

WO0019982

Title:

SPRAY COATED MICROPARTICLES FOR USE IN NEEDLELESS SYRINGES

Abstract:

Spray-coated pharmaceutical powder compositions for transdermal administration using a needleless syringe comprise seed particles coated with a pharmaceutical composition, the said coated seed particles having an average size of about 10 to 100 μm and having an envelope density ranging from about 0.1 to about 25g/cm³.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 9/16	A1	(11) International Publication Number: WO 00/19982 (43) International Publication Date: 13 April 2000 (13.04.00)
(21) International Application Number: PCT/IB99/01859 (22) International Filing Date: 1 October 1999 (01.10.99) (30) Priority Data: 60/102,726 1 October 1998 (01.10.98) US (71) Applicant: POWDERJECT RESEARCH LIMITED [GB/GB]; 4 Robert Robinson Avenue, Oxford Science Park, Oxford OX4 4GA (GB). (72) Inventor: MAA, Yuh-Fun; 651 Sequoia Avenue, Millbrae, CA 94030 (US). (74) Agent: WOODS, Geoffrey, Corlett; J.A. Kemp & Co., 14 South Square, Gray's Inn, London WC1R 5LX (GB).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: SPRAY COATED MICROPARTICLES FOR USE IN NEEDLELESS SYRINGES (57) Abstract <p>Spray-coated pharmaceutical powder compositions for transdermal administration using a needleless syringe comprise seed particles coated with a pharmaceutical composition, the said coated seed particles having an average size of about 10 to 100 μm and having an envelope density ranging from about 0.1 to about 25g/cm³.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakistan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

SPRAY COATED MICROPARTICLES FOR USE IN NEEDLELESS SYRINGES

Technical Field

5 The invention relates to a method for producing powdered pharmaceutical compositions. More specifically, the invention relates to a method for forming dense, substantially solid particles from pharmaceutical compositions, where the particulate compositions are particularly suitable for transdermal particle delivery from a needleless syringe system.

10 Background

The ability to deliver pharmaceuticals through skin surfaces (transdermal delivery) provides many advantages over oral or parenteral delivery techniques. In particular, transdermal delivery provides a safe, convenient and noninvasive alternative to traditional drug administration systems, conveniently avoiding the major problems associated with oral delivery (e.g., variable rates of absorption and metabolism, gastrointestinal irritation and/or bitter or unpleasant drug tastes) or parenteral delivery (e.g., needle pain, the risk of introducing infection to treated individuals, the risk of contamination or infection of health care workers caused by accidental needle-sticks and the disposal of used needles).

20 However, despite its clear advantages, transdermal delivery presents a number of its own inherent logistical problems. The passive delivery of drugs through intact skin necessarily entails the transport of molecules through a number of structurally different tissues, including the stratum corneum, the viable epidermis, the papillary dermis, and the capillary walls in order for the drug to gain entry into the blood or lymph system. Transdermal delivery systems must therefore be able to overcome the various resistances presented by each type of tissue. In light of the above, a number of alternatives to passive transdermal delivery have been developed. These alternatives include the use of skin penetration enhancing agents, or "permeation enhancers," to increase skin permeability, as well as non-chemical modes such as the use of iontophoresis, electroporation or ultrasound. However, these alternative techniques often give rise to their own unique side effects, such as skin irritation or sensitization. Thus,

the spectrum of pharmaceuticals that can be safely and effectively administered using traditional transdermal delivery methods has remained limited.

More recently, a novel transdermal drug delivery system that entails the use of a needleless syringe to fire powders (i.e., solid drug-containing particles) in controlled doses into and through intact skin has been described. In particular, commonly owned U.S. Patent No. 5,630,796 to Bellhouse et al. describes a needleless syringe that delivers pharmaceutical particles entrained in a supersonic gas flow. The needleless syringe is used for transdermal delivery of powdered drug compounds and compositions, for delivery of genetic material into living cells (e.g., gene therapy) and for the delivery of biopharmaceuticals to skin, muscle, blood or lymph. The needleless syringe can also be used in conjunction with surgery to deliver drugs and biologics to organ surfaces, solid tumors and/or to surgical cavities (e.g., tumor beds or cavities after tumor resection). In theory, practically any pharmaceutical agent that can be prepared in a substantially solid, particulate form can be safely and easily delivered using such devices.

One particular needleless syringe generally comprises an elongate tubular nozzle having a rupturable membrane initially closing the passage through the nozzle and arranged substantially adjacent to the upstream end of the nozzle. Particles of a therapeutic agent to be delivered are disposed adjacent to the rupturable membrane and are delivered using an energizing means which applies a gaseous pressure to the upstream side of the membrane sufficient to burst the membrane and produce a supersonic gas flow (containing the pharmaceutical particles) through the nozzle for delivery from the downstream end thereof. The particles can thus be delivered from the needleless syringe at delivery velocities of between Mach 1 and Mach 8 which are readily obtainable upon the bursting of the rupturable membrane.

Another needleless syringe configuration generally includes the same elements as described above, except that instead of having the pharmaceutical particles entrained within a supersonic gas flow, the downstream end of the nozzle is provided with a bistable diaphragm which is moveable between a resting "inverted" position (in which the diaphragm presents a concavity on the downstream face to contain the pharmaceutical particles) and an active "everted"

position (in which the diaphragm is outwardly convex on the downstream face as a result of a supersonic shockwave having been applied to the upstream face of the diaphragm). In this manner, the pharmaceutical particles contained within the concavity of the diaphragm are expelled at a supersonic initial velocity from the device for transdermal delivery thereof to a targeted skin or mucosal surface.

Transdermal delivery using either of the above-described needleless syringe configurations is carried out with particles having an approximate size that generally ranges between 0.1 and 250 μm . For drug delivery, an optimal particle size is usually at least about 10 to 15 μm (the size of a typical cell). For gene delivery, an optimal particle size is generally substantially smaller than 10 μm . Particles larger than about 250 μm can also be delivered from the device, with the upper limitation being the point at which the size of the particles would cause untoward damage to the skin cells. The actual distance which the delivered particles will penetrate depends upon particle size (e.g., the nominal particle diameter assuming a roughly spherical particle geometry), particle density, the initial velocity at which the particle impacts the skin surface, and the density and kinematic viscosity of the skin. In this regard, optimal particle densities for use in needleless injection generally range between about 0.1 and 25 g/cm^3 , preferably between about 0.8 and 1.5 g/cm^3 , and injection velocities generally range between about 100 and 3,000 m/sec .

Summary of the Invention

It is a primary object of the invention to provide a spray-coated powder composition for administration from a needleless syringe. It is also a primary object of the invention to provide suitable spray-coating methods for producing such powder compositions.

In one aspect of the invention, a spray-coated powder composition for administration from a needleless syringe is provided. The powder composition is formed from seed particles that are coated with an aqueous pharmaceutical composition. More especially, the spray-coated powder composition comprises seed particles coated with a pharmaceutical composition, the said coated seed particles having an average size of about 10 to 100 μm and having an envelope

density ranging from about 0.1 to about 25 g/cm³.

The coated seed particles can have an average size of about 20 to 70 μm. Preferably, they have an envelope density ranging from about 0.8 to about 1.5 g/cm³. The coated seed particles typically have a substantially spherical
5 aerodynamic shape and/or a substantially uniform, nonporous surface. The powders may also be characterized in that the coated seed particles have a pharmaceutical composition loading of about 1 to 50 wt%. The spray-coated powder compositions can contain, as the active pharmaceutical agent, any small molecule drug substance, organic or inorganic chemical, vaccine, or peptide
10 (polypeptide and/or protein) of interest.

In another aspect of the invention, a method for preparing the spray-coated powder composition is provided. The method comprises spray-coating an aqueous pharmaceutical composition onto seed particles under conditions sufficient to provide coated particles having an average size of about 10 to 100
15 μm and an envelope density ranging from about 0.1 to about 25 g/cm³. In one particular embodiment, the method entails the steps of: (a) suspending the seed particles in a reaction chamber using a hot air flow; (b) atomizing an aqueous pharmaceutical composition into a fine spray and introducing the spray into the reaction chamber; (c) allowing the spray to spread over the surface of the
20 suspended seed particles to coat them with a thin film; and then (d) drying the coated seed particles. If desired, the aqueous pharmaceutical composition can be sprayed into the reaction chamber in a direction that is transverse to the direction of the hot air flow.

It is an advantage of the invention that free-flowing powder compositions
25 can be produced having well defined particle size, density and mechanical properties which collectively are suitable for transdermal delivery from a needleless syringe. Further advantages of the methods of the invention include flexible pharmaceutical loading (up to about 50 wt%), overall process efficiency (no need for post-formulation fractionation, classification or sieving operations),
30 and the methods are readily scalable. The invention further provides:

- a dosage receptacle for a needleless syringe, said receptacle containing a therapeutically effective amount of a spray-coated powder composition of

the invention; and

- a needleless syringe which is loaded with such a powder composition.

These and other objects, aspects, embodiments and advantages of the present invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

Detailed Description of the Preferred Embodiments

Before describing the present invention in detail, it is to be understood that this invention is not limited to particularly exemplified compositions or process parameters as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments of the invention only, and is not intended to be limiting.

All publications, patents and patent applications cited herein, whether *supra* or *infra*, are hereby incorporated by reference in their entirety.

It must be noted that, as used in this specification and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to “a particle” includes a mixture of two or more such particles, reference to “an excipient” includes mixtures of two or more such excipients, and the like.

A. Definitions

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. Although a number of methods and materials similar or equivalent to those described herein can be used in the practice of the present invention, the preferred materials and methods are described herein.

In describing the present invention, the following terms will be employed, and are intended to be defined as indicated below.

By “transdermal delivery,” applicant intends to include both transdermal (“percutaneous”) and transmucosal routes of administration, i.e., delivery by passage of a drug or pharmaceutical agent through the skin or mucosal tissue. *See, e.g., Transdermal Drug Delivery: Developmental Issues*

and Research Initiatives, Hadgraft and Guy (eds.), Marcel Dekker, Inc., (1989); *Controlled Drug Delivery: Fundamentals and Applications*, Robinson and Lee (eds.), Marcel Dekker Inc., (1987); and *Transdermal Delivery of Drugs*, Vols. 1-3, Kydonieus and Berner (eds.), CRC Press, (1987).

- 5 As used herein, the term “pharmaceutical” or “pharmaceutical agent” intends any compound or composition of matter which, when administered to an organism (human or animal) induces a desired pharmacologic and/or physiologic effect by local and/or systemic action. The term therefore encompasses those compounds or chemicals traditionally regarded as drugs, as well as
- 10 biopharmaceuticals including molecules such as peptides, hormones, nucleic acids, gene constructs and the like. More particularly, the term “pharmaceutical” or “pharmaceutical agent” includes compounds or compositions for use in all of the major therapeutic areas including, but not limited to, anti-infectives such as antibiotics and antiviral agents; analgesics and analgesic combinations; local and
- 15 general anesthetics; anorexics; antiarthritics; antiasthmatic agents; anticonvulsants; antidepressants; antihistamines; anti-inflammatory agents; antinauseants; antineoplastics; antipruritics; antipsychotics; antipyretics; antispasmodics; cardiovascular preparations (including calcium channel blockers, beta-blockers, beta-agonists and antiarrhythmics); antihypertensives; diuretics;
- 20 vasodilators; central nervous system stimulants; cough and cold preparations; decongestants; diagnostics; hormones; bone growth stimulants and bone resorption inhibitors; immunosuppressives; muscle relaxants; psychostimulants; sedatives; tranquilizers; therapeutic proteins (e.g., antigens, antibodies, growth factors, cytokines, interleukins, lymphokines, interferons, enzymes, etc.),
- 25 peptides and fragments thereof (whether naturally occurring, chemically synthesized or recombinantly produced); and nucleic acid molecules (polymeric forms of two or more nucleotides, either ribonucleotides (RNA) or deoxyribonucleotides (DNA) including both double- and single-stranded molecules, gene constructs, expression vectors, antisense molecules and the like).
- 30 The above pharmaceuticals or pharmaceutical agents, alone or in combination with other agents, are typically prepared as pharmaceutical compositions which can contain one or more added materials such as carriers,

vehicles, and/or excipients. "Carriers," "vehicles" and "excipients" generally refer to substantially inert materials which are nontoxic and do not interact with other components of the composition in a deleterious manner. These materials can be used to increase the amount of solids in particulate pharmaceutical compositions. Examples of suitable carriers include water, silicone, gelatin, waxes, and like materials. Examples of normally employed "excipients," include pharmaceutical grades of carbohydrates including monosaccharides, disaccharides, cyclodextrans, and polysaccharides (e.g., dextrose, sucrose, lactose, trehalose, raffinose, mannitol, sorbitol, inositol, dextrans, and maltodextrans); starch; cellulose; salts (e.g. sodium or calcium phosphates, calcium sulfate, magnesium sulfate); citric acid; tartaric acid; glycine; high molecular weight polyethylene glycols (PEG); Pluronic; surfactants; and combinations thereof. Generally, when carriers and/or excipients are used, they are used in amounts ranging from about 0.1 to 99 wt% of the pharmaceutical composition.

The terms "individual" and "subject" are used interchangeably herein to refer to any member of the subphylum cordata, including, without limitation, humans and other primates, including non-human primates such as chimpanzees and other apes and monkey species; farm animals such as cattle, sheep, pigs, goats and horses; domestic mammals such as dogs and cats; laboratory animals including rodents such as mice, rats and guinea pigs; birds, including domestic, wild and game birds such as chickens, turkeys and other gallinaceous birds, ducks, geese, and the like. The terms do not denote a particular age. Thus, both adult and newborn individuals are intended to be covered. The methods described herein are intended for use in any of the above vertebrate species, since the immune systems of all of these vertebrates operate similarly.

The term "powder," as used herein, refers to a composition that consists of substantially solid particles that can be delivered transdermally using a needleless syringe device. The particles that make up the powder can be characterized on the basis of a number of parameters including, but not limited to, the average particle size, the average particle density, particle morphology (e.g., particle aerodynamic shape and particle surface characteristics), and particle

penetration energy (P.E.).

The average particle size of the powders produced according to the present invention can vary widely and is generally between about 10 and 100 μm , more typically between about 20 and 70 μm . The average particle size of the powder can be measured as a mass mean aerodynamic diameter (MMAD) using conventional techniques such as microscopic techniques (where particles are sized directly and individually rather than grouped statistically), absorption of gasses, permeability or time of flight. If desired, automatic particle-size counters can be used (e.g., Aerosizer Counter, Coulter Counter, HIAC Counter, or Gelman Automatic Particle Counter) to ascertain the average particle size.

Actual particle density, or "absolute density," can be readily ascertained using known quantification techniques such as helium pycnometry and the like. Alternatively, envelope ("tap") density measurements can be used to assess the density of a particulate pharmaceutical composition produced according to the methods of the invention. Envelope density information is particularly useful in characterizing the density of objects of irregular size and shape. Envelope density is the mass of an object divided by its volume, where the volume includes that of its pores and small cavities but excludes interstitial space. A number of methods of determining envelope density are known in the art, including wax immersion, mercury displacement, water absorption and apparent specific gravity techniques. A number of suitable devices are also available for determining envelope density, for example, the GeoPycTM Model 1360, available from the Micromeritics Instrument Corp. The difference between the absolute density and envelope density of a sample pharmaceutical composition provides information about the sample's percentage total porosity and specific pore volume.

Particle morphology, particularly the aerodynamic shape of a particle, can be readily assessed using standard light microscopy. It is preferred that the particles which make up the instant powders have a substantially spherical or at least substantially elliptical aerodynamic shape. It is also preferred that the particles have an axis ratio of 3 or less to avoid the presence of rod- or needle-shaped particles. These same microscopic techniques can also be used to assess the particle surface characteristics, e.g., the amount and extent of surface voids or

degree of porosity.

Particle penetration energies can be ascertained using a number of conventional techniques, for example a metallized film P.E. test.

5 B. General Methods

 In one embodiment, a powdered pharmaceutical composition is provided, wherein the powder composition (which is comprised of particles) is produced using a spray-coating technique. The powders are suitable for transdermal administration from a needleless syringe delivery system, and as such, the
10 particles which make up the powdered composition must have sufficient physical strength to withstand sudden acceleration to several times the speed of sound and the impact with, and passage through, the skin and tissue. The particles are formed by spray-coating an aqueous pharmaceutical composition onto suitable seed particles. The particles can be delivered from a needleless syringe system
15 such as those described in commonly owned International Publication Nos. WO 94/24263, WO 96/04947, WO 96/12513, and WO 96/20022, all of which are incorporated herein by reference.

 The aqueous pharmaceutical compositions can contain, as the active pharmaceutical agent, any small molecule drug substance, organic or inorganic
20 chemical, vaccine, or peptide (polypeptide and/or protein) of interest. In particular embodiments, the pharmaceutical agent is a biopharmaceutical preparation of a peptide, polypeptide, protein or any other such biological molecule. Exemplary peptide and protein formulations include, without limitation, insulin; calcitonin; octreotide; endorphin; liprecin; pituitary hormones
25 (e.g., human growth hormone and recombinant human growth hormone (hGH and rhGH), HMG, desmopressin acetate, etc); follicle luteoids; growth factors (such as growth factor releasing factor (GFRF), somatostatin, somatotropin and platelet-derived growth factor); asparaginase; chorionic gonadotropin; corticotropin (ACTH); erythropoietin (EPO); epoprostenol (platelet aggregation
30 inhibitor); glucagon; interferons; interleukins; menotropins (urofollitropin, which contains follicle-stimulating hormone (FSH); and luteinizing hormone (LH)); oxytocin; streptokinase; tissue plasminogen activator (TPA); urokinase;

vasopressin; desmopressin; ACTH analogues; angiotensin II antagonists; antidiuretic hormone agonists; bradykinin antagonists; CD4 molecules; antibody molecules and antibody fragments (e.g., Fab, Fab₂, Fv and sFv molecules); IGF-1; neurotrophic factors; colony stimulating factors; parathyroid hormone and
5 agonists; parathyroid hormone antagonists; prostaglandin antagonists; protein C; protein S; renin inhibitors; thrombolytics; tumor necrosis factor (TNF); vaccines (particularly peptide vaccines including subunit and synthetic peptide preparations); vasopressin antagonists analogues; and α -1 antitrypsin. Additionally, nucleic acid preparation, such as vectors or gene constructs for use
10 in subsequent gene delivery, can be used.

The pharmaceutical agent is typically prepared as an aqueous pharmaceutical composition using a suitable aqueous carrier, along with suitable excipients, protectants, solvents, salts, surfactants, buffering agents and the like. Suitable excipients can include free-flowing particulate solids that do not thicken
15 or polymerize upon contact with water, which are innocuous when administered to an individual, and do not significantly interact with the pharmaceutical agent in a manner that alters its pharmaceutical activity. In general, excipients which are sticky, or have high hygroscopicity are avoided particularly for powder formulations where the pharmaceutical is loaded onto the seed particle at a high
20 concentration (e.g., >10 wt%). Examples of normally employed excipients include, but are not limited to, pharmaceutical grades of dextrose, sucrose, lactose, trehalose, mannitol, sorbitol, inositol, dextran, starch, cellulose, sodium or calcium phosphates, calcium carbonate, calcium sulfate, sodium citrate, citric acid, tartaric acid, glycine, high molecular weight polyethylene glycols (PEG),
25 and combinations thereof. Suitable solvents include, but are not limited to, methylene chloride, acetone, methanol, ethanol, isopropanol and water. Generally pharmaceutically acceptable salts having molarities ranging from about 1 mM to 2M can be used. Pharmaceutically acceptable salts include, for example, mineral acid salts such as hydrochlorides, hydrobromides, phosphates,
30 sulfates, and the like; and the salts of organic acids such as acetates, propionates, malonates, benzoates, and the like. A thorough discussion of pharmaceutically acceptable excipients, vehicles and auxiliary substances is available in

REMINGTON'S PHARMACEUTICAL SCIENCES (Mack Pub. Co., N.J. 1991), incorporated herein by reference.

The seed particles can be comprised of any parenterally acceptable powder (e.g., crystalline or amorphous), are selected to have good flowability
5 (i.e., are fluidizable), and are sufficiently dense for efficient use with needleless transdermal delivery systems. Crystalline particles are generally preferred due to their inherently high particle density and overall penetration energy. Seed particles having an overall spherical or at least elliptical shape are preferred. Particles generally are selected to have an axis ratio of 3 or less, for example 2 or
10 less or 1.5 or less, in order to avoid rod- or needle-shaped particles which are difficult to reprocess and are generally less flowable.

Suitable seed particles can be comprised of any pharmaceutically acceptable carbohydrate (e.g., sugars such as lactose, mannitol, trehalose, etc.), polysaccharide, starch, biodegradable polymer (e.g. PLGA, a copolymer of L-
15 lactic acid and glycolic acid), or the like. The seed particles can have an average size of about 5 to 100 μm , for example about 10 to 95 μm or about 20 to 70 μm . Seed particle preparations having a substantially homogenous average particle size can be readily obtaining using standard sieving or other particle classification methodologies.

20 The spray-coated powders can be formed using any standard spray-coating processing apparatus. In this regard, batch-type fluid-bed processors have long been used to perform drying, granulation, and coating operations in the pharmaceutical industry for preparing solid dosage forms. Olsen, K.W. (1989) "Batch fluid-bed processing equipment: A design overview," Part I., *Pharm. Technol.* 13:34-46, Olsen, K.W. (1989) "Batch fluid-bed processing equipment:
25 A design overview," Part II., *Pharm. Technol.* 13:39-50. With the advent of the Würster spray coater, seed particles as small as 50 μm in size can, at least in theory, be coated. Iyer et al. (1993) *Drug Devel. Ind. Pharm.* 19:981-989. However, to date, the spray coating of seed particles having an average size of
30 100 μm or less has been limited, particularly for protein or peptide pharmaceuticals. Maa et al. (1996) *Intl. J. Pharmaceutics* 144:47-59.

Spray coating processors that can fluidize seed particles of 10 μm or

larger, for example 20 μm and larger, and which can atomize a fine spray (droplet size of 30 μm or less, preferably 10 μm or less) are preferred. Suitable processors include any commercially available Würster spray coater, or Würster HS spray coater (available from Glatt Air Techniques, Inc.). For fluid-bed processing, the
5 spray coating processor can utilize any suitable spraying method which is selected in consideration of the desired characteristics for the finished product. These spraying methods (e.g., top, bottom or tangential (rotary coater)) are generally known to those skilled in the art.

The liquid delivery system for the spray coat processor typically utilizes a
10 binary nozzle, where the aqueous pharmaceutical composition is supplied at a relatively low pressure through an orifice and is atomized by air. Pneumatic nozzles can be used to produce smaller droplets. The atomization conditions, including atomization gas flow rate, atomization gas pressure, liquid flow rate, etc., can be controlled to produce droplets from the pharmaceutical composition
15 having an average diameter of about 30 μm or less, with droplets having an average size of 10 μm or less being preferred. Typically, the atomizing air pressure, liquid flow rate and the fluidizing air temperature and volume are the most significant process variables and have the greatest effect over the particle characteristics of the resultant coated particles. Drying temperature conditions of
20 about 50-150 °C inlet temperature and about 30-100 °C outlet temperature are preferred. The thickness of the pharmaceutical coat can be controlled by the drying time, and the present methods can provide spray-coated powders formed from seed particles loaded with from about 1 to 50 wt% (e.g., about 0.5 to 15 wt% of active pharmaceutical agent in compositions containing both active
25 pharmaceutical agent and carrier), preferably >10 wt% of the aqueous pharmaceutical composition.

If desired, a secondary coating process can be used to provide further structural integrity in the coated particles, for example, where the spray-coated powder particles are coated with a standard sugar excipient using the same sort of
30 spray-coating procedure as described herein above. In some cases, it may be desirable to coat the spray-coated powder particles with the same sugar used as the seed (e.g., mannitol, lactose, trehalose, or the like). Other secondary coating

materials include, but are not limited to, pharmaceutical grades of carbohydrates including monosaccharides, disaccharides, cyclodextrans, and polysaccharides (e.g., dextrose, sucrose, raffinose, mannose, sorbitol, inositol, dextrans, and maltodextrans); starch; cellulose; salts (e.g. sodium or calcium phosphates, calcium sulfate, magnesium sulfate); citric acid; tartaric acid; glycine; high molecular weight polyethylene glycols (PEG); Pluronics; surfactants; and combinations thereof. The secondary coating material can also be used to optimize the particles for delivery to mucosal target surfaces (e.g., by coating the spray-coated powder particles with a lipid), or to alter or retard solubility characteristics of the particles after delivery into an aqueous environment (e.g., by applying a secondary coating containing a salt, starch, dextran, or the like).

The spray-coating methods of the present invention can be used to produce powders that are suitable for transdermal delivery from a needleless syringe delivery device. Typical powders are characterized in that the individual particles have an average size in the range of about 20 to 70 μm , an envelope density ranging from about 0.1 to about 25 g/cm^3 , preferably ranging from about 0.8 to about 1.5 g/cm^3 , and have a substantially spherical aerodynamic shape with a substantially uniform, nonporous surface.

The particles which make up the spray-coated powders of the present invention will also have a particle penetration energy suitable for transdermal delivery from a needleless syringe device. Such penetration energies can conveniently be assessed using a metallized film P.E. measuring procedure as follows. A metallized film material (e.g., a 125 μm polyester film having a 350 Å layer of aluminum deposited on a single side) is used as a substrate into which the powder is fired from a needleless syringe (e.g., the needleless syringe described in U.S. Patent No. 5,630,796 to Bellhouse et al.) at an initial velocity of about 100 to 3000 m/sec. The metallized film is placed, with the metal coated side facing upwards, on a suitable surface. A needleless syringe loaded with a spray-coated powder produced according to the methods of the invention is placed with its spacer contacting the film, and then fired. Residual powder is removed from the metallized film surface using a suitable solvent. Penetration energy is then assessed using a BioRad Model GS-700 imaging densitometer to

scan the metallized film, and a personal computer with a SCSI interface and loaded with MultiAnalyst software (BioRad) and Matlab software (Release 5.1, The MathWorks, Inc.) is used to assess the densitometer reading. A program is used to process the densitometer scans made using either the transmittance or
5 reflectance method of the densitometer. The penetration energy of the spray-coated powders should be equivalent to, or better than that of reprocessed mannitol particles of the same size (mannitol particles that are freeze-dried, compressed, ground and sieved according to the methods of commonly owned International Publication No. WO 97/48485, incorporated herein by reference).

10 Once produced, the spray-coated powders of the present invention can be packaged in individual unit dosages. As used herein, a "unit dosage" intends a dosage receptacle containing a therapeutically effective amount of a spray-coated pharmaceutical produced according to the methods of the present invention. The dosage receptacle is generally one which fits within a needleless syringe device to
15 allow for transdermal delivery from the device. Such receptacles can be capsules, foil pouches, sachets, cassettes, or the like.

C. Experimental

Below are examples of specific embodiments for carrying out the methods
20 of the present invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some experimental error and deviation should, of course, be allowed for.

25

EXAMPLE 1

The following spray-coated powder formulations are made using the methods of the present invention.

30

Formulation 1:

Seed particles: 500 grams of lactose (Pharmatose, 100M & 200M,

Crompton & Knowle), sieved to provide an average particle size of 20-75 μm by jet sieve.

Aqueous pharmaceutical composition: Lysozyme (50%) and trehalose (50%) at a total solid concentration of 20%.

- 5 Spray coater: GPCG-1 (Glatt Air), operated at the following coating conditions: air inlet temperature = 85°C, air outlet temperature = 42°C, liquid feed = 15 mL/min, air velocity in the bed = 3.5 m/sec., coating loading 10% of lysozyme, and a coating time = 35 min.

10 Formulation 2:

Seed particles: 300 grams of lactose (Pharmatose, 100M & 200M, Crompton & Knowle), sieved to provide an average particle size of 20-75 μm by jet sieve.

- 15 Aqueous pharmaceutical composition: s-Calcitonin (20%), mannitol (30%), and trehalose (50%) at a total solid concentration of 10%.

Spray coater: Precision coater (MP-1, Niro), operated at the following coating conditions: air inlet temperature = 68°C, air outlet temperature = 34°C, liquid feed = 12 mL/min, air velocity in the bed = 3.5 m/sec., coating loading 5% of s-Calcitonin, and a coating time = 63 min.

20

Formulation 3:

Seed particles: 300 grams of mannitol (Merck), sieved to provide an average particle size of 20-75 μm by jet sieve.

- 25 Aqueous pharmaceutical composition: recombinant human growth hormone (rhGH) (50%), mannitol (20%), glycine (10%), and trehalose (20%) at a total solid concentration of 10%.

- 30 Spray coater: Precision coater (MP-1, Niro), operated at the following coating conditions: air inlet temperature = 75°C, air outlet temperature = 40°C, liquid feed = 13 mL/min, air velocity in the bed = 3.5 m/sec., coating loading 5% of rhGH, and a coating time = 24 min.

Formulation 4:

Seed particles: 500 grams of mannitol (Merck), sieved to provide an average particle size of 20-75 μm by jet sieve.

5 Aqueous pharmaceutical composition: Bovine serum albumin (BSA) (100%) at a total solid concentration of 10%.

Spray coater: GPCG-1 (Glatt Air), operated at the following coating conditions: air inlet temperature = 85°C, air outlet temperature = 42°C, liquid feed = 15 mL/min, air velocity in the bed = 3.5 m/sec., coating loading 15% of BSA, and a coating time = 50 min.

10

Formulation 5:

Seed particles: 300 grams of lactose (Pharmatose, 100M & 200M, Crompton & Knowle), sieved to provide an average particle size of 20-75 μm by jet sieve.

15 Aqueous pharmaceutical composition: Diphtheria toxoids vaccine (DPT) (2%), mannitol (30%), glycine (8%), and trehalose (60%) at a total solid concentration of 10%.

Spray coater: Precision coater (MP-1, Niro), operated at the following coating conditions: air inlet temperature = 75°C, air outlet temperature = 40°C, 20 liquid feed = 13 mL/min, air velocity in the bed = 3.5 m/sec., coating loading 0.5% of DPT, and a coating time = 58 min.

The spray-coated powder formulation of each Example is suitable for transdermal administration from a needleless syringe and is characterized in that 25 the individual spray-coated particles have an average size in the range of about 20 to 70 μm , an envelope density ranging from about 0.1 to about 25 g/cm³, preferably ranging from about 0.8 to about 1.5 g/cm³, and have a substantially spherical aerodynamic shape with a substantially uniform, nonporous surface.

Accordingly, novel spray-coated powder compositions, and methods for 30 producing these compositions have been described. Although preferred embodiments of the subject invention have been described in some detail, it is understood that obvious variations can be made without departing from the spirit

and the scope of the invention as defined by the appended claims.

What is claimed is:

1. A spray-coated powder composition for administration from a
needleless syringe, said powder composition comprising seed particles coated
5 with a pharmaceutical composition, the said coated seed particles having an
average size of about 10 to 100 μm and having an envelope density ranging from
about 0.1 to about 25 g/cm^3 .
2. The powder composition of claim 1, wherein said seed particles
10 are crystalline particles.
3. The powder composition of claim 1 or 2, wherein the seed
particles have an axis ratio of 3 or less.
4. The powder composition of any one of the preceding claims,
15 wherein the seed particles are selected from the group consisting of lactose,
mannitol, trehalose, polysaccharides, starches, and biodegradable polymers.
5. The powder composition of any one of the preceding claims,
20 wherein said coated seed particles have an average size of about 20 to 70 μm .
6. The powder composition of any one of the preceding claims,
wherein said coated seed particles have an envelope density ranging from about
0.8 to about 1.5 g/cm^3 .
25
7. The powder composition of any one of the preceding claims,
wherein said coated seed particles have a substantially spherical aerodynamic
shape.
8. The powder composition of any one of the preceding claims,
30 wherein said coated seed particles have a substantially uniform, nonporous
surface.

9. The powder composition of any one of the preceding claims, wherein said coated seed particles have a pharmaceutical composition loading of about 1 to 50 wt%.

5 10. A method for preparing a spray-coated powder for administration from a needleless syringe, said method comprising spray-coating an aqueous pharmaceutical composition onto seed particles under conditions sufficient to provide coated particles having an average size of about 10 to 100 μm and an envelope density ranging from about 0.1 to about 25 g/cm^3 .

10

11. The method of claim 10 comprising the steps of:

(a) suspending the seed particles in a reaction chamber using a heated air flow;

(b) atomizing an aqueous pharmaceutical composition into a spray
15 and introducing said spray into the reaction chamber;

(c) allowing the spray to spread over the surface of said suspended seed particles to coat said seed particles with a film; and

(d) drying the coated seed particles.

20 12. The method of claim 11, wherein the aqueous pharmaceutical composition is sprayed into the reaction chamber in a direction that is transverse to the direction of the heated air flow.

25 13. A dosage receptacle for a needleless syringe, said receptacle containing a therapeutically effective amount of a spray-coated powder composition as claimed in any one of claims 1 to 9.

14. The receptacle of claim 13, which is selected from the group consisting of capsules, foil pouches, sachets and cassettes.

30

15. A needleless syringe which is loaded with a spray-coated powder composition as claimed in any one of claims 1 to 9.

INTERNATIONAL SEARCH REPORT

Intern. Application No
PCT/IB 99/01859

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K9/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94 23738 A (MEDISORB TECHNOLOGIES INTERNATIONAL) 27 October 1994 (1994-10-27) the whole document	1-15
A	WO 97 48485 A (POWDERJECT RESEARCH) 24 December 1997 (1997-12-24) the whole document	1-15
A	WO 93 00991 A (SMITHKLINE BEECHAM) 21 January 1993 (1993-01-21) the whole document	1-15
A	EP 0 532 777 A (HAARMANN & REIMER) 24 March 1993 (1993-03-24) the whole document	1-15
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"G" document member of the same patent family

Date of the actual completion of the international search

9 March 2000

Date of mailing of the international search report

16/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Ventura Amat, A

INTERNATIONAL SEARCH REPORT

Intern. Application No
PCT/IB 99/01859

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 12513 A (OXFORD BIOSCIENCES) 2 May 1996 (1996-05-02) cited in the application the whole document ----	1-15
A	WO 96 20022 A (BELLHOUSE) 4 July 1996 (1996-07-04) cited in the application the whole document ----	1-15
A	WO 96 04947 A (OXFORD BIOSCIENCES) 22 February 1996 (1996-02-22) cited in the application the whole document ----	1-15
A	WO 94 24263 A (OXFORD BIOSCIENCES) 27 October 1994 (1994-10-27) cited in the application the whole document ----	1-15
A	WO 98 10750 A (POWDERJECT RESEARCH) 19 March 1998 (1998-03-19) the whole document -----	1-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 99/01859

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9423738 A	27-10-1994	AU 6707194 A CA 2160878 A EP 0696200 A JP 8510639 T NZ 265818 A	08-11-1994 27-10-1994 14-02-1996 12-11-1996 22-09-1997
WO 9748485 A	24-12-1997	AU 3102597 A CA 2258554 A EP 0912239 A	07-01-1998 24-12-1997 06-05-1999
WO 9300991 A	21-01-1993	AT 157559 T AU 659192 B AU 2166492 A DE 69222043 D DE 69222043 T DK 593552 T EP 0593552 A ES 2106875 T GR 3025311 T HK 1002628 A IE 79672 B JP 6508779 T ZA 9204882 A	15-09-1997 11-05-1995 11-02-1993 09-10-1997 05-02-1998 27-10-1997 27-04-1994 16-11-1997 27-02-1998 04-09-1998 20-05-1998 06-10-1994 27-05-1993
EP 532777 A	24-03-1993	NONE	
WO 9612513 A	02-05-1996	AU 3703895 A CA 2199417 A DE 69507321 D DE 69507321 T EP 0788386 A JP 10507390 T	15-05-1996 02-05-1996 25-02-1999 15-07-1999 13-08-1997 21-07-1998
WO 9620022 A	04-07-1996	AT 183657 T AU 699463 B AU 4271496 A BG 101600 A BR 9510464 A CA 2208590 A CN 1171055 A CZ 9701937 A DE 69511723 D EP 0799064 A FI 972554 A HU 77064 A JP 10512467 T NO 972897 A NZ 297542 A PL 320919 A SK 81797 A	15-09-1999 03-12-1998 19-07-1996 27-02-1998 29-06-1999 04-07-1996 21-01-1998 12-11-1997 30-09-1999 08-10-1997 16-06-1997 02-03-1998 02-12-1998 20-06-1997 28-10-1999 10-11-1997 04-03-1998
WO 9604947 A	22-02-1996	AU 3229095 A CA 2197711 A DE 69506559 D DE 69506559 T EP 0776224 A JP 10504214 T	07-03-1996 22-02-1996 21-01-1999 06-05-1999 04-06-1997 28-04-1998

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 99/01859

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9424263 A	27-10-1994	AT 148497 T	15-02-1997
		AU 674742 B	09-01-1997
		AU 6435194 A	08-11-1994
		BG 61993 B	30-12-1998
		BG 100047 A	30-04-1996
		BR 9406455 A	02-01-1996
		CA 2159452 A	27-10-1994
		CN 1120852 A	17-04-1996
		CZ 9502608 A	15-05-1996
		DE 69401651 D	13-03-1997
		DE 69401651 T	15-05-1997
		DK 693119 T	28-07-1997
		EP 0693119 A	24-01-1996
		EP 0734737 A	02-10-1996
		EP 0951917 A	27-10-1999
		ES 2098131 T	16-04-1997
		FI 954788 A	06-10-1995
		GR 3022939 T	30-06-1997
		HK 1000351 A	06-03-1998
		HU 73516 A, B	28-08-1996
		JP 8509604 T	15-10-1996
		LV 11833 A	20-08-1997
		LV 11833 B	20-12-1997
		NO 953994 A	06-10-1995
		NZ 263606 A	22-08-1997
		PL 311005 A	22-01-1996
		SG 48696 A	18-05-1998
		SI 693119 T	31-10-1997
		SK 124895 A	08-01-1997
		US 5630796 A	20-05-1997
		ZA 9402442 A	10-04-1995
WO 9810750 A	19-03-1998	AU 4307997 A	02-04-1998
		EP 0928190 A	14-07-1999